arXiv:1410.6416v1 [math.CA] 5 Oct 2014

ON THE MAXIMAL OPERATORS OF VILENKIN-FEJÉR MEANS

GEORGE TEPHNADZE

Abstract. The main aim of this paper is to prove that the maximal operator $\widetilde{\sigma}^* f := \sup_{n\in P} \frac{|\sigma_n f|}{\log^2(n+1)}$ is bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$, where $\sigma_n f$ is Fejér means of bounded Vilenkin-Fourier series.

2000 Mathematics Subject Classification. 42C10.
Key words and phrases: Vilenkin system, Fejér means, martingale Hardy space.

1. Introduction

In one-dimensional case the weak type inequality

$$\mu\left(\sigma^* f > \lambda\right) \leq \frac{c}{\lambda} \|f\|_1, \qquad (\lambda > 0)$$

can be found in Zygmund [19] for the trigonometric series, in Schipp [11] for Walsh series and in Pál, Simon [10] for bounded Vilenkin series. Again in one-dimensional, Fujji [4] and Simon [13] verified that σ^* is bounded from H_1 to L_1. Weisz [16] generalized this result and proved the boundedness of σ^* from the martingale space H_p to the space L_p for $p > 1/2$. Simon [12] gave a counterexample, which shows that boundedness does not hold for $0 < p < 1/2$. The counterexample for $p = 1/2$ due to Goginava [7], (see also [3]). In the endpoint case $p = 1/2$ two positive results was showed. Weisz [18] proved that σ^* is bounded from the Hardy space $H_{1/2}$ to the space weak-$L_{1/2}$. For Walsh-Paley system in 2008 Goginava [6] proved that the maximal operator $\widetilde{\sigma}^*$ defined by

$$\widetilde{\sigma}^* f := \sup_{n\in P} \frac{|\sigma_n f|}{\log^2(n+1)}$$

is bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$. He also proved that for any nondecreasing function $\varphi : P_+ \to [1, \infty)$ satisfying the condition

$$\overline{\lim_{n\to\infty}} \frac{\log^2(n+1)}{\varphi(n)} = +\infty \tag{1}$$

the maximal operator

$$\sup_{n\in P} \frac{|\sigma_n f|}{\varphi(n)}$$

is not bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$.

For Walsh-Kaczmarz system analogical theorem is proved in [9].

The main aim of this paper is to prove that the maximal operator $\widetilde{\sigma}^{*}f$ with respect to Vilenkin system is bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$ (see Theorem 1).

We also prove that under the condition (1) the maximal operator

$$\sup_{n\in P}\frac{|\sigma_n f|}{\varphi\left(n\right)}$$

is not bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$. Actually, we prove stronger result (see Theorem 2) than the unboundedness of the maximal operator $\widetilde{\sigma}^{*}f$ from the Hardy space $H_{1/2}$ to the spaces $L_{1/2}$. In particular, we prove that

$$\sup_{n\in P}\left\|\frac{\sigma_n f}{\varphi\left(n\right)}\right\|_{L_{1/2}}=\infty.$$

2. Definitions and Notation

Let P_{+} denote the set of the positive integers , $P:=P_{+}\cup\{0\}$.

Let $m:=(m_{0,}m_{1,...})$ denote a sequence of the positive integers not less than 2.

Denote by

$$Z_{m_k}:=\{0,1,...m_k-1\}$$

the additive group of integers modulo m_k.

Define the group G_m as the complete direct product of the group Z_{m_j} with the product of the discrete topologies of Z_{m_j} 's.

The direct product μ of the measures

$$\mu_k\left(\{j\}\right):=1/m_k,\qquad\left(j\in Z_{m_k}\right),$$

is the Haar measure on G_m , with $\mu\left(G_m\right)=1$.

If $\sup\limits_{n}m_n<\infty$, then we call G_m a bounded Vilenkin group. If the generating sequence m is not bounded then G_m is said to be an unbounded Vilenkin group. **In this paper we discuss bounded Vilenkin groups only.**

The elements of G_m represented by sequences

$$x:=(x_0,x_{1,...},x_{j,...}),\qquad\left(\ x_k\in Z_{m_k}\right).$$

It is easy to give a base for the neighborhood of G_m :

$$I_0\left(x\right):=G_m,$$

$$I_n(x):=\{y\in G_m\mid y_0=x_0,...y_{n-1}=x_{n-1}\},\ \left(x\in G_m,\ n\in P\right).$$

Denote $I_n:=I_n\left(0\right)$, for $n\in P$ and $\overline{I_n}:=G_m\setminus I_n$.

Let

$$e_n:=(0,...,0,x_n=1,0,...)\in G_m,\qquad\left(n\in P\right).$$

Denote

$$I_N^{k,\,l} := \begin{cases} I_N(0,...,0,x_k \neq 0,0,...,0,x_l \neq 0,x_{l+1,...,}\, x_{N-1}\,),\ k<l<N, \\ I_N(0,...,0,x_k \neq 0,0,...,,0),\ l=N. \end{cases}$$

and

$$I_N^{k,\alpha,l,\beta} := I_N(0,...,0,x_k=\alpha,0,...,0,x_l=\beta,x_{l+1,...,}\, x_{N-1}\,),\ k<l<N.$$

It is evident

$$I_N^{k,\,l} = \bigcup_{\alpha=1}^{m_k-1} \bigcup_{\beta=1}^{m_l-1} I_N^{k,\alpha,l,\beta} \tag{2}$$

and

$$\bar{I_N} = \left(\bigcup_{k=0}^{N-2} \bigcup_{l=k+1}^{N-1} I_N^{k,\,l} \right) \cup \left(\bigcup_{k=0}^{N-1} I_N^{k,\,N} \right). \tag{3}$$

If we define the so-called generalized number system based on m in the following way :

$$M_0 := 1, \qquad M_{k+1} := m_k M_k\,, \qquad (k \in P).$$

then every $n \in P$ can be uniquely expressed as $n = \sum_{k=0}^{\infty} n_j M_j$, where $n_j \in Z_{m_j}$ $(j \in P)$ and only a finite number of n_j's differ from zero. Let $|n| := \max \{j \in P,\, n_j \neq 0\}$.

Denote by $L_1\,(G_m)$ the usual (one dimensional) Lebesque space.

Next, we introduce on G_m an ortonormal system which is called the Vilenkin system.

At first define the complex valued function $r_k\,(x) : G_m \to C$, the generalized Rademacher functions as

$$r_k\,(x) := \exp\left(2\pi i x_k/m_k\right), \qquad \left(i^2=-1,\ x \in G_m,\ k \in P\right).$$

Now define the Vilenkin system $\psi := (\psi_n : n \in P)$ on G_m as:

$$\psi_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}\,(x)\,, \qquad (n \in P)\,.$$

Specifically, we call this system the Walsh-Paley one if m=2.

The Vilenkin system is ortonormal and complete in $L_2\,(G_m)$ [1, 14].

Now we introduce analogues of the usual definitions in Fourier-analysis.

If $f \in L_1\,(G_m)$ we can establish the the Fourier coefficients, the partial sums of the Fourier series, the Fejér means, the Dirichlet and Fejér kernels

with respect to the Vilenkin system ψ in the usual manner:

$$\begin{aligned}
\widehat{f}(k) &:= \int_{G_m} f\overline{\psi}_k d\mu, \qquad (\, k \in P\,),\\
S_n f &:= \sum_{k=0}^{n-1} \widehat{f}(k)\,\psi_k, \qquad (\, n \in P_+,\ S_0 f := 0),\\
\sigma_n f &:= \frac{1}{n}\sum_{k=0}^{n-1} S_k f, \qquad (\, n \in P_+\,),\\
D_n &:= \sum_{k=0}^{n-1} \psi_k\,, \qquad (\, n \in P_+\,),\\
K_n &:= \frac{1}{n}\sum_{k=0}^{n-1} D_k, \qquad (\, n \in P_+\,),
\end{aligned}$$

Recall that

$$D_{M_n}(x) = \begin{cases} M_n, & \text{if } x \in I_n,\\ 0, & \text{if } x \notin I_n. \end{cases} \tag{4}$$

It is well-known that

$$\sup_n \int_{G_m} |K_n(x)|\, d\mu(x) \le c < \infty, \tag{5}$$

$$n\,|K_n(x)| \le c\sum_{A=0}^{|n|} M_A\,|K_{M_A}(x)|\,. \tag{6}$$

The norm (or quasinorm) of the space $L_p(G_m)$ is defined by

$$\|f\|_p := \left(\int_{G_m} |f(x)|^p\, d\mu(x)\right)^{1/p}, \qquad (0 < p < \infty)\,.$$

The $\sigma-$algebra generated by the intervals $\{I_n(x) : x \in G_m\}$ will be denoted by F_n $(n \in P)$. Denote by $f = \left(f^{(n)}, n \in P\right)$ a martingale with respect to F_n $(n \in P)$. (for details see e.g. [15]). The maximal function of a martingale f is defend by

$$f^* = \sup_{n\in P}\left|f^{(n)}\right|.$$

In case $f \in L_1$, the maximal functions are also be given by

$$f^*(x) = \sup_{n\in P}\frac{1}{|I_n(x)|}\left|\int_{I_n(x)} f(u)\,\mu(u)\right|.$$

For $0 < p < \infty$, the Hardy martingale spaces $H_p\,(G_m)$ consist of all martingales for which

$$\|f\|_{H_p} := \|f^*\|_{L_p} < \infty.$$

If $f \in L_1$,then it is easy to show that the sequence $(S_{M_n}(f) : n \in P)$ is a martingale. If $f = \left(f^{(n)}, n \in P\right)$ is martingale then the Vilenkin-Fourier coefficients must be defined in a slightly different manner:

$$\widehat{f}(i) := \lim_{k\to\infty} \int_{G_m} f^{(k)}(x)\, \overline{\Psi}_i(x)\, d\mu(x).$$

The Vilenkin-Fourier coefficients of $f \in L_1(G_m)$ are the same as those of the martingale $(S_{M_n}(f) : n \in P)$ obtained from f .

For the martingale f we consider maximal operators

$$\begin{aligned} \sigma^* f &= \sup_{n\in P} |\sigma_n f|, \\ \widetilde{\sigma}^* f &: = \sup_{n\in P} \frac{|\sigma_n f|}{\log^2(n+1)}. \end{aligned}$$

A bounded measurable function a is p-atom, if there exist a dyadic interval I, such that

$$\begin{cases} a) & \int_I a d\mu = 0, \\ b) & \|a\|_\infty \le \mu(I)^{-1/p}, \\ c) & \operatorname{supp}(a) \subset I. \end{cases}$$

3. Formulation of Main Results

Theorem 1. *The maximal operator*

$$\widetilde{\sigma}^* f := \sup_{n\in P} \frac{|\sigma_n f|}{\log^2(n+1)}$$

is bounded from the Hardy space $H_{1/2}(G_m)$ to the space $L_{1/2}(G_m)$.

Theorem 2. *Let $\varphi : P_+ \to [1, \infty)$ be a nondecreasing function satisfying the condition*

$$\overline{\lim_{n\to\infty}} \frac{\log^2(n+1)}{\varphi(n)} = +\infty.$$

Then there exists a martingale $f \in H_{1/2}$, such that

$$\sup_{n\in P} \left\| \frac{\sigma_n f}{\varphi(n)} \right\|_{L_{1/2}} = \infty.$$

Corollary 1. *Under condition (1) the maximal operator*

$$\sup_{n\in P} \frac{|\sigma_n f|}{\varphi(n)}$$

is not bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$.

4. AUXILIARY PROPOSITIONS

Lemma 1. [17] *Suppose that an operator T is sublinear and for some $0 < p \leq 1$*

$$\int_{\overline{I}} |Ta|^p \, d\mu \leq c_p < \infty$$

for every p-atom a, where I denote the support of the atom. If T is bounded from L_∞ to L_∞, then

$$\|Tf\|_{L_p(G_m)} \leq c_p \|f\|_{H_p(G_m)}.$$

Lemma 2. [2, 8] *Let $2 < A \in P_+$, $k \leq s < A$ and $q_A = M_{2A} + M_{2A-2} + ... + M_2 + M_0$. Then*

$$q_{A-1} \left| K_{q_{A-1}}(x) \right| \geq \frac{M_{2k} M_{2s}}{4},$$

for

$$x \in I_{2A} \left(0, ..., x_{2k} \neq 0, 0, ..., 0, x_{2s} \neq 0, x_{2s+1}, ...x_{2A-1}\right),$$

$$k = 0, \ 1, ..., \ A-3, \qquad s = k+2, \ k+3, ..., \ A-1.$$

Lemma 3. [5] *Let $A > t$, $t, A \in P$, $z \in I_t \setminus I_{t+1}$. Then*

$$K_{M_A}(z) = \begin{cases} 0, & \text{if } z - z_t e_t \notin I_A, \\ \frac{M_t}{1 - r_t(z)}, & \text{if } z - z_t e_t \in I_A. \end{cases}$$

Lemma 4. *Let $x \in I_N^{k,l}$, $\qquad k = 0, ..., N-1$, $l = k+1, ..., N$. Then*

$$\int_{I_N} |K_n(x-t)| \, d\mu(t) \leq \frac{c M_l M_k}{M_N^2}, \quad \text{when } \ n \geq M_N.$$

Proof. Let $x \in I_N^{k,\alpha,l,\beta}$. Then applying lemma 3 we have

$$K_{M_A}(x) = 0, \ \text{when } \ A > l.$$

Hence we can suppose that $A \leq l$.

Let $k < A \leq l$. Then we have

$$|K_{M_A}(x)| = \frac{M_k}{|1 - \ r_k(x)|} \leq \frac{m_k M_k}{2\pi \ \alpha}. \tag{7}$$

Let $x \in I_N^{k,\,l}$, $0 \leq k < l \leq N-1$ and $t \in I_N$. Since $x - t \in I_N^{k,\,l}$ and $n \geq M_N$ combining (2) (6) and (7) we obtain

$$n\left|K_n\left(x\right)\right| \leq c\sum_{A=0}^{l-1} M_A M_k \leq cM_k M_l \tag{8}$$

and

$$\int_{I_N} \left|K_n\left(x-t\right)\right| d\mu\left(t\right) \leq \frac{cM_k M_l}{M_N^2}. \tag{9}$$

Let $x \in I_N^{k,N}$, then applying (6) we have

$$\int_{I_N} n\left|K_n\left(x-t\right)\right| d\mu\left(t\right) \leq \sum_{A=0}^{|n|} M_A \int_{I_N} \left|K_{M_A}\left(x-t\right)\right| d\mu\left(t\right). \tag{10}$$

Let

$$\begin{cases} x = \left(0,...,0,x_k \neq 0,0,...0,x_N,x_{N+1},x_q,...,x_{|n|-1},,...\right), \\ t = \left(0,...,0,x_N,...x_{q-1},t_q \neq x_q,t_{q+1},...,t_{|n|-1},...\right),\ q = N,...,|n|-1. \end{cases}$$

Using Lemma 3 in (10) it is easy to show that

$$\begin{aligned} & \int_{I_N} \left|K_n\left(x-t\right)\right| d\mu\left(t\right) \\ \leq\ & \frac{c}{n}\sum_{A=0}^{q-1} M_A \int_{I_N} M_k d\mu\left(t\right) \\ \leq\ & \frac{cM_k M_q}{nM_N} \leq \frac{cM_k}{M_N}. \end{aligned} \tag{11}$$

Let

$$\begin{cases} x = \left(0,...,0,x_m \neq 0,0,...,0,x_N,x_{N+1},x_q,...,x_{|n|-1},...\right), \\ t = \left(0,0,...,x_N,...,x_{|n|-1},t_{|n|},...\right). \end{cases}.$$

If we apply Lemma 3 in (10) we obtain

$$\begin{aligned} & \int_{I_N} \left|K_n\left(x-t\right)\right| d\mu\left(t\right) \\ \leq\ & \frac{c}{n}\sum_{A=0}^{|n|-1} M_A \int_{I_N} M_k d\mu\left(t\right) \leq \frac{cM_k}{M_N}. \end{aligned} \tag{12}$$

Combining (9), (11) and (12) we complete the proof of lemma 4.

5. Proof of the Theorems

Proof of Theorem 1. By Lemma 1, the proof of theorem 1 will be complete, if we show that

$$\int_{\overline{I}_N} \left(\sup_{n\in P} \frac{|\sigma_n a|}{\log^2 (n+1)} \right)^{1/2} d\mu \leq c < \infty,$$

for every $1/2$-atom a, where I denotes the support of the atom and bounded from L_∞ to L_∞. The boundedness follows from (5).

Let a be an arbitrary $1/2$-atom with support I and $\mu(I) = M_N^{-1}$. We may assume that $I = I_N$. It is easy to see that $\sigma_n(a) = 0$ when $n \leq M_N$. Therefore we can suppose that $n > M_N$.

Since $\|a\|_\infty \leq cM_N^2$ we can write

$$\begin{aligned} & \frac{|\sigma_n(a)|}{\log^2 (n+1)} \\ \leq & \frac{1}{\log^2 (n+1)} \int_{I_N} |a(t)|\, |K_n(x-t)|\, d\mu(t) \\ \leq & \frac{\|a\|_\infty}{\log^2 (n+1)} \int_{I_N} |K_n(x-t)|\, d\mu(t) \\ \leq & \frac{cM_N^2}{\log^2 (n+1)} \int_{I_N} |K_n(x-t)|\, d\mu(t). \end{aligned}$$

Let $x \in I_N^{k,\,l}$, $0 \leq k < l \leq N$. Then from Lemma 4 we get

$$\frac{|\sigma_n(a)|}{\log^2 (n+1)} \leq \frac{cM_N^2}{N^2} \frac{M_l M_k}{M_N^2} = \frac{cM_l M_k}{N^2}. \tag{13}$$

Combining (3) and (13) we obtain

$$\begin{aligned}
&\int_{\overline{I_N}} |\widetilde{\sigma}^{*} a\left(x\right)|^{1/2} d\mu\left(x\right) \\
&= \sum_{k=0}^{N-2}\sum_{l=k+1}^{N-1}\sum_{x_j=0, j\in\{l+1,\ldots,N-1\}}^{m_j-1} \int_{I_N^{k,\,l}} \left|\widetilde{\sigma}^{*} a\left(x\right)\right|^{1/2} d\mu\left(x\right) \\
&\quad +\sum_{k=0}^{N-1}\int_{I_N^{k,N}} |\widetilde{\sigma}^{*} a\left(x\right)|^{1/2} d\mu\left(x\right) \\
&\leq c\sum_{k=0}^{N-2}\sum_{l=k+1}^{N-1}\frac{m_{l+1}\ldots m_{N-1}}{M_N}\frac{\sqrt{M_l M_k}}{N} \\
&\quad +c\sum_{k=0}^{N-1}\frac{1}{M_N}\frac{\sqrt{M_N M_k}}{N} \\
&\leq c\sum_{k=0}^{N-2}\sum_{l=k+1}^{N-1}\frac{\sqrt{M_k}}{N\sqrt{M_l}} + c\sum_{k=0}^{N-1}\frac{1}{\sqrt{M_N}}\frac{\sqrt{M_k}}{N} \leq c < \infty.
\end{aligned}$$

Which complete the proof of Theorem 1.

Proof of Theorem 2. Let $\{\lambda_k; k\in P_+\}$ be an increasing sequence of the positive integers such that

$$\lim_{k\to\infty}\frac{\log^2\left(\lambda_k\right)}{\varphi\left(\lambda_k\right)} = \infty.$$

It is evident that for every λ_k there exists a positive integers $m_k^{'}$ such that $q_{m_k^{'}} \leq \lambda_k < q_{m_k^{'}+1} < c\, q_{m_k^{'}}$. Since $\varphi\left(n\right)$ is nondecreasing function.We have

$$\overline{\lim_{k\to\infty}}\frac{\left(m_k^{'}\right)^2}{\varphi\left(q_{m_k^{'}}\right)} \geq c\lim_{k\to\infty}\frac{\log^2\left(\lambda_k\right)}{\varphi\left(\lambda_k\right)} = \infty. \tag{14}$$

Let $\{n_k; k\in P_+\}\subset\{m_k^{\prime}; k\in P_+\}$ such that

$$\lim_{k\to\infty}\frac{n_k^2}{\varphi\left(q_{n_k}\right)} = \infty$$

and

$$f_{n_k}\left(x\right) = D_{M_{2n_k+1}}\left(x\right) - D_{M_{2n_k}}\left(x\right).$$

It is evident

$$\widehat{f}_{n_k}\left(i\right) = \begin{cases} 1, \text{ if } i = M_{2n_k},\ldots,M_{2n_k+1}-1, \\ 0, \text{otherwise.}\end{cases}$$

Then we can write

$$S_i\left(f_{n_k}(x)\right)=\begin{cases} D_i\left(x\right)-D_{M_{2n_k}}\left(x\right)\ , & \text{if } i=M_{2n_k},...,M_{2n_k+1}-1,\\ f_{n_k}\left(x\right)\ , & \text{if } i\geq M_{2n_k+1},\\ 0, & \text{otherwise.}\end{cases} \tag{15}$$

From (4) we get

$$\begin{aligned} &\|f_{n_k}\|_{H_{1/\,2}} \\ &= \left\|\sup_{n\in P}S_{M_n}f_{n_k}\right\|_{L_{1/2}} \\ &= \left\|D_{M_{2n_k+1}}-D_{M_{2n_k}}\right\|_{L_{1/2}} \\ &= \left(\int_{I_{2n_k}\setminus I_{2n_k+1}}M_{2n_k}^{1/2}d\mu\left(x\right)+\int_{I_{2n_k+1}}\left(M_{2n_k+1}-M_{2n_k}\right)^{1/2}d\mu\left(x\right)\right)^2 \\ &= \left(\frac{m_{2n_k}-1}{M_{2n_k+1}}M_{2n_k}^{1/2}+\frac{\left(m_{2n_k}-1\right)^{1/2}}{M_{2n_k+1}}M_{2n_k}^{1/2}\right)^2 \\ &\leq \frac{c}{M_{2n_k}}. \end{aligned} \tag{16}$$

By (15) we can write:

$$\begin{aligned} &\frac{\left|\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)\right|}{\varphi\left(q_{n_k}\right)} \\ &= \frac{1}{\varphi\left(q_{n_k}\right)q_{n_k}}\left|\sum_{j=0}^{q_{n_k}-1}S_jf_{n_k}(x)\right| \\ &= \frac{1}{\varphi\left(q_{n_k}\right)q_{n_k}}\left|\sum_{j=M_{2n_k}}^{q_{n_k}-1}S_jf_{n_k}(x)\right| \\ &= \frac{1}{\varphi\left(q_{n_k}\right)q_{n_k}}\left|\sum_{j=M_{2n_k}}^{q_{n_k}-1}\left(D_j\left(x\right)-D_{M_{2n_k}}\left(x\right)\right)\right| \\ &= \frac{1}{\varphi\left(q_{n_k}\right)q_{n_k}}\left|\sum_{j=0}^{q_{n_k-1}-1}\left(D_{j+M_{2n_k}}\left(x\right)-D_{M_{2n_k}}\left(x\right)\right)\right| \end{aligned}$$

Since

$$D_{j+M_{2n_k}}\left(x\right)-D_{M_{2n_k}}\left(x\right)=\psi_{M_{2n_k}}D_j,\ \ j=1,2,..,M_{2n_k}-1,$$

we obtain

$$\begin{aligned}
&\frac{\left|\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)\right|}{\varphi\left(q_{n_k}\right)}\\
&= \frac{1}{\varphi\left(q_{n_k}\right)q_{n_k}}\left|\sum_{j=0}^{q_{n_k-1}-1} D_j\left(x\right)\right|\\
&= \frac{1}{\varphi\left(q_{n_k}\right)}\frac{q_{n_k-1}}{q_{n_k}}\left|K_{q_{n_k-1}}\left(x\right)\right|.
\end{aligned}$$

Let $x \in I_{2n_k}^{2s,2l}$. Then from Lemma 2 we obtain

$$\frac{\left|\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)\right|}{\varphi\left(q_{n_k}\right)} \geq \frac{cM_{2s}M_{2l}}{M_{2n_k}\varphi\left(q_{n_k}\right)}.$$

Hence we can write:

$$\begin{aligned}
&\int_{G_m}\left|\frac{\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)}{\varphi\left(q_{n_k}\right)}\right|^{1/2} d\mu\left(x\right)\\
&\geq \sum_{s=0}^{n_k-3}\sum_{l=s+1}^{n_k-1}\sum_{x_{2l+1}=0}^{m_{2l+1}}\cdots\sum_{x_{2n_k-1}=0}^{m_{2n_k-1}}\int_{I_{2n_k}^{2s,2l}}\left|\frac{\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)}{\varphi\left(q_{n_k}\right)}\right|^{1/2} d\mu\left(x\right)\\
&\geq c\sum_{s=0}^{n_k-3}\sum_{l=s+1}^{n_k-1}\frac{m_{2l+1}\ldots m_{2n_k-1}}{M_{2n_k}}\frac{\sqrt{M_{2s}M_{2l}}}{\sqrt{\varphi\left(q_{n_k}\right)M_{2n_k}}}\\
&\geq c\sum_{s=0}^{n_k-3}\sum_{l=s+1}^{n_k-1}\frac{\sqrt{M_{2s}}}{\sqrt{M_{2l}M_{2n_k}\varphi\left(q_{n_k}\right)}}\\
&\geq \frac{cn_k}{\sqrt{M_{2n_k}\varphi\left(q_{n_k}\right)}}.
\end{aligned}$$

From (16) we have

$$\begin{aligned}
&\frac{\left(\int_{G_m}\left|\frac{\sigma_{q_{n_k}}\left(f_{n_k}(x)\right)}{\varphi(n_k)}\right|^{1/2} d\mu\left(x\right)\right)^2}{\left\|f_{n_k}\right\|_{H_{1/\,2}}}\\
&\geq \frac{cn_k^2}{M_{2n_k}\varphi\left(q_{n_k}\right)}M_{2n_k}\\
&\geq \frac{cn_k^2}{\varphi\left(q_{n_k}\right)} \to \infty, \qquad \text{when} \qquad k \to \infty.
\end{aligned}$$

Theorem 2 is proved.

References

[1] Agaev, G. N., Vilenkin, N.Ya., Dzhafarly G.M., Rubinshtein A. I.: Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehim, 1981 (in Russian).

[2] Blahota, I., Gát, G., Goginava, U.: Maximal operators of Fejer means of double Vilenkin-Fourier series, Colloq. Math. 107 (2007), no. 2, 287–296.

[3] Blahota, I., Gát, G., Goginava, U.: Maximal operators of Fejér means of Vilenkin-Fourier series. JIPAM. J. Inequal. Pure Appl. Math. 7 (2006), 1- 7 .

[4] Fujji, N. J.: A maximal inequality for H_1 functions on the generalized Walsh-Paley group, Proc. Amer. Math. Soc. 77 (1979), lll-116.

[5] Gát, G.: Cesàro means of integrable functions with respect to unbounded Vilenkin systems. J. Approx. Theory 124 (2003), no. 1, 25–43.

[6] Goginava, U.: Maximal operators of Fejér-Walsh means. Acta Sci. Math. (Szeged) 74 (2008), no. 3-4, 615–624.

[7] Goginava, U.: The maximal operator of the Fejér means of the character system of the p-series field in the Kaczmarz rearrangement. Publ. Math. Debrecen 71 (2007), no. 1-2, 43–55.

[8] Goginava, U.: Maximal operators of Fejér means of double Walsh-Fourier series. Acta Math. Hungar. 115 (2007), no. 4, 333–340.

[9] Goginava, U., Nagy, K.: On the maximal operator of Walsh-Kaczmarz-Fejer means, Czechoslovak Math. J. (to appear).

[10] Pál, J., Simon, J.: On a generalization of the comncept of derivate, Acta Math. Hung., 29 (1977), 155-164.

[11] Schipp, F.: Certain rearranngements of series in the Walsh series, Mat. Zametki, 18 (1975), 193-201.

[12] Simon, P.: Cesàro summability wish respect to two-parameter Walsh sistems, Monatsh. Math .,131 (2000),321-334.

[13] Simon, P.: Inverstigations with respect to the Vilenkin sistem, Annales Univ. Sci. Budapest Eotv., Sect. Math., 28 (1985) 87-101.

[14] Vilenkin, N.Ya.: A class of complate ortonormal systems, Izv. Akad. Nauk. U.S.S.R., Ser. Mat., 11 (1947), 363-400.

[15] Weisz, F.: Martingale Hardy spaces and their applications in Fourier Analysis, Springer, Berlin-Heideiberg-New York, 1994.

[16] Weisz, F.: Cesáro summability of one and two-dimensional Fourier series, Anal. Math. 5 (1996), 353-367. Appl. Math. 7 (2006), 1- 7 .

[17] Weisz, F.: Summability of multi-dimensional Fourier series and Hardy space, Kluwer Academic, Dordrecht, 2002.

[18] Weisz, F.: Weak type inequalities for the Walsh and bounded Ciesielski systems. Anal. Math. 30 (2004), no. 2, 147–160.

[19] Zygmund, A.: Trigonometric Series, Vol. 1, Cambridge Univ. Press, 1959.

G. Tephnadze, department of Mathematics, Faculty of Exact and Natural Sciences, Tbilisi State University, Chavchavadze str. 1, Tbilisi 0128, Georgia

E-mail address: giorgitephnadze@gmail.com